

# A 2D Bose Gas to Study Quantum Hydrodynamic Instabilities

P H C CUNHA<sup>1</sup>, G T BELUMAT<sup>1</sup>, G M NOVAES<sup>1</sup>, L B PIETRO<sup>1</sup>, K M FARIAS<sup>1</sup>, AND P C M CASTILHO<sup>1</sup>

<sup>1</sup>*Grupo de optica, University of São Paulo, São Carlos Institute of Physics, São Carlos, Brazil*  
Contact Email: patricia.castilho@ifsc.usp.br

Hydrodynamic instabilities in classical fluids, often characterized by the exponential growth of a well-defined pattern, are encountered in many mundane situations. Increasing our understanding of the underlying mechanisms of such instabilities has a strong impact in modeling fluids dynamics in a broad scenario and in fundamental problems such as the transition from laminar to turbulent flow. In the context of quantum fluids, quantum hydrodynamic instabilities (QHI) are related to the superfluid properties of these systems offering a new approach when studying superfluidity. Thanks to their high degree of control and simple detection techniques, ultracold atomic gases are ideal platforms to engender and observe such QHI. In this work, we present the design of a new experimental setup capable to address the specific conditions for the onset of different hydrodynamic instabilities in a 2D Bose gas. Combining the flexibility of the optical potentials created with the use of a DMD with the capability of tuning the atomic interactions offered by potassium-39, we aim to initially observe the onset of the quantum Rayleigh-Taylor instability and follow its dynamical evolution.